## Day Three

#### **Instructions:**

- 1. Use this page for **day three of a three-day diary**. The three days you choose do not have to be in a row; however, each day must be a full 24 hours during which it is convenient for you to measure every void.
- 2. Begin recording when you wake up in the morning and continue for 24 hours.
- 3. Make a separate entry each time you void, leak or have anything to drink.
- 4. Measure voids (mL measurements) using the hat. If you are not at home, use the following scale to measure the amount of urine voided: 1=small, 2=medium, 3=large.
- 5. Measure fluid intake in ounces.
- 6. When recording a leak, use the following scale: 1=drops/damp, 2=wet/soaked, 3=bladder emptied. Indicate your activity during the leak and if you had an urge to void (yes or no; strong or not)

Time	Fluid Intake (Type and amount in ounces)	Amount Voided (mL)	Involuntary Urine Leakage (scale 1-3)	Activity During Leak	Was there an urge?
Example					
7:05		350 mL			
7:40	Coffee 8 oz				
8.10			2	Walking from car	Strongurge

Example Diary

#### Day 3 Date:

Time	Fluid Intake (Type and amount in ounces)	Amount Voided (mL)	Involuntary Urine Leakage (scale 1-3)	Activity During Leak	Was there an urge?

Any other notations you care to make:



### **Kegel Exercises**

#### **Kegel Exercises Strengthen Pelvic Muscles**

If you need additional help identifying and exercising your pelvic muscles, we have a physical therapist who is trained to evaluate your muscle strength and help you learn how to perform Kegel exercises. Ask your provider for information.

#### **Find Your Pelvic Muscles**

- Try to stop your urinary stream. If you succeed, you have identified the right muscles to exercise.
- Exercising the muscles around the rectum will also strengthen those around the vagina and under the bladder. Squeeze the muscles that would prevent gas from escaping from your rectum.
- Do not tighten your stomach and back muscles or squeeze your legs together as you isolate and contract your pelvic muscles.

#### **Kegel Exercise Routine**

- Begin your routine lying down. As you get stronger, you can exercise while sitting or standing.
- Pull in and hold a pelvic muscle squeeze for five seconds; relax for five seconds.
- Do 10 sets of repetitions three times a day.
- Increase your hold by one second each week until you are contracting for 10 seconds.
- Rest and breathe between contractions.

Source: American Urogynecologic Society

# Premier Health Premier Physician Network

#### **Lifestages Centers for Women**

**(**937) 277-8988

BeavercreekHuber Heights4235 Indian Ripple Rd.6251 Miami Valley WaySuite 210Suite 130Beavercreek, Ohio 45440Huber Heights, Ohio 45424

CentervilleMiddletown6611 Clyo Road200 Medical Center DriveSuite DSuite 160

Centerville, Ohio 45459 Middletown, Ohio 45005

EnglewoodSpringboro9000 North Main Street630 N. Main StreetSuite 232Suite 200Englewood, Ohio 45415Springboro, Ohio 45066

S-C-WMN93579-6/2019 • LSBD-2018

## **Bladder Diary**

One- to Three-Days

Lifestages Centers for Women





LifestagesCentersForWomen.com



## Day One

#### **Instructions:**

- 1. If your provider has asked for a **one-day diary**, use this page. If your provider has asked for a three-day diary, begin with this page. The days you choose do not have to be in a row.
- 2. Choose a 24-hour period when it is convenient for you to measure every void.
- 3. Begin recording when you wake up in the morning and continue for a full 24 hours. (see example, below)
- 4. Make a separate entry each time you void, leak or have anything to drink.
- 5. Measure voids (mL measurements) using the hat you were given in our office. If you are not at home, use the following scale to measure the amount of urine voided: 1=small, 2=medium, 3=large.
- 6. Measure fluid intake in ounces.
- 7. When recording a leak, use the following scale: 1=drops/damp, 2=wet/soaked, 3=bladder emptied. Indicate your activity during the leak and if you had an urge to void (yes or no; strong or not)

Time	Fluid Intake (Type and amount in ounces)	Amount Voided (mL)	Involuntary Urine Leakage (scale 1-3)	Activity During Leak	Was there an urge?
Example					
7:05		350 mL			
7:40	Coffee 8 oz				
8-10			2	Walking from car	Strong urge

Example Diary

#### Day 1 Date:

Time	Fluid Intake (Type and amount in ounces)	Amount Voided (mL)	Involuntary Urine Leakage (scale 1-3)	Activity During Leak	Was there an urge?

Any other notations you care to make:



## Day Two

#### **Instructions:**

- 1. Use this page for **day two of a three-day diary**. The days you choose do not have to be in a row; however, each day must be a full 24 hours during which it is convenient for you to measure every void.
- 2. Begin recording when you wake up in the morning and continue for 24 hours.
- 3. Make a separate entry each time you void, leak or have anything to drink.
- 4. Measure voids (mL measurements) using the hat. If you are not at home, use the following scale to measure the amount of urine voided: 1=small, 2=medium, 3=large.
- 5. Measure fluid intake in ounces.
- 6. When recording a leak, use the following scale: 1=drops/damp, 2=wet/soaked, 3=bladder emptied. Indicate your activity during the leak and if you had an urge to void (yes or no; strong or not)

Time	Fluid Intake (Type and amount in ounces)	Amount Voided (mL)	Involuntary Urine Leakage (scale 1-3)	Activity During Leak	Was there an urge?	Ex
Example						
7:05		350 mL				D
7:40	Coffee 8 oz					1
8:10			2	Walking from car	Strong urge	

Example Diary

#### Day 2 Date:

<u>-</u>					
Time	Fluid Intake (Type and amount in ounces)	Amount Voided (mL)	Involuntary Urine Leakage (scale 1-3)	Activity During Leak	Was there an urge?

Any other notations you care to make:



